

40 C.F.R. § 180.1

Definitions and interpretations.

- (a) *Administrator*, without qualification, means the Administrator of the Environmental Protection Agency.
- (b) *Agency*, without qualification, means the Environmental Protection Agency.
- (c) *FFDCA* means the Federal Food, Drug, and Cosmetic Act, as amended, 21 U.S.C. 301-392.
- (d) Raw agricultural commodities include, among other things, fresh fruits, whether or not they have been washed and colored or otherwise treated in their unpeeled natural form; vegetables in their raw or natural state, whether or not they have been stripped of their outer leaves, waxed, prepared into fresh green salads, etc.; grains, nuts, eggs, raw milk, meats, and similar agricultural produce. It does not include foods that have been processed, fabricated, or manufactured by cooking, freezing, dehydrating, or milling.
- (e) Where a raw agricultural commodity bearing a pesticide chemical residue that has been exempted from the requirement of a tolerance, or which is within a tolerance permitted under FFDCA section 408, is used in preparing a processed food, the processed food will not be considered unsafe within the meaning of FFDCA sections 402 and 408(a), despite the lack of a tolerance or exemption for the pesticide chemical residue in the processed food, if:
 - (1) The pesticide chemical has been used in or on the raw agricultural commodity in conformity with a tolerance under this section;
 - (2) The pesticide chemical residue has been removed to the extent possible in good manufacturing practice; and
 - (3) The concentration of the pesticide chemical residue in the processed food is not greater than the tolerance prescribed for the pesticide chemical residue on the raw agricultural commodity.
- (f) For the purpose of computing fees as required by § 180.33, each group of related crops listed in § 180.34(e) and each crop group or subgroup listed in § 180.41 is counted as a single raw agricultural commodity in a petition or request for tolerances or exemption from the requirement of a tolerance.
- (g) Tolerances and exemptions established for pesticide chemicals in or on the general category of raw agricultural commodities listed in column A apply to the corresponding specific raw agricultural commodities listed in column B. However, a tolerance or exemption for a specific commodity in column B does not apply to the general category in column A.

A	B
Alfalfa	<i>Medicago sativa</i> L. <i>Subsp. sativa</i> , (alfalfa, lucerne); <i>Onobrychis viciifolia</i> Scop. (sainfoin, holy clover, esparcet); and <i>Lotus corniculatus</i> L. (trefoil); and varieties and/or hybrids of these.

Banana	Banana, plantain.
Basil (<i>Ocimum</i> spp.)	Basil (<i>Ocimum basilicum</i> L.); Basil, American (<i>Ocimum americanum</i> L.); Basil, Greek (<i>Ocimum minimum</i> L.); Basil, holy (<i>Ocimum tenuiflorum</i> L.); Basil, lemon (<i>Ocimum x citriodorum</i> Vis.); Basil, Russian (<i>Ocimum gratissimum</i> L.)
Bean	<i>Cicer arietinum</i> (chickpea, garbanzo bean); <i>Lupinus</i> spp. (including, but not limited to, Andean lupin, blue lupin, grain lupin, sweet lupin, white sweet lupin, white lupin, and yellow lupin). <i>Phaseolus</i> spp. (including, but not limited to, black bean, cranberry bean, dry bean, field bean, French bean, garden bean, great northern bean, green bean, kidney bean, lima bean, navy bean, pink bean, pinto bean, red bean, scarlet runner bean, snap bean, tepary bean, yellow bean, and wax bean); Broad bean (fava bean, faba bean); Goa bean (asparagus pea and winged bean); <i>Vigna</i> spp. (including adzuki bean, asparagus bean, blackeyed pea, catjang bean, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, and yardlong bean); Guar bean; Horse gram; Jackbean; Lablab bean (hyacinth bean); Morama bean; African yam bean; American potato bean; Vegetable soybean (edamame); Sword bean; Velvetbean; Winged pea; cultivars, varieties and/or hybrids of these commodities. [Note: A variety of pesticide tolerances have been previously established for pea and/or bean. Chickpea/garbanzo bean is also listed in the definition for “pea”. For garbanzo bean/chickpea only, the highest established pea or bean tolerance will apply to pesticide residues found in this commodity].
Bean, dry, seed	All beans in the entry “Bean” in dry seed form.
Bean, edible podded	All beans in the entry “Bean” in edible podded form.
Bean, succulent	All beans in the entry “Bean” in edible podded or succulent shelled form.
Bean, succulent shelled	All beans in the entry “Bean” in succulent shelled form.
Blackberry	<i>Rubus eubatus</i> (including bingleberry, black satin berry, boysenberry Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, Lavacaberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangerberry, ravenberry, rossberry, Shawnee blackberry, and varieties and/or hybrids of these).
Broccoli	Broccoli, Chinese broccoli (gai lon, white flowering broccoli).
Cabbage	Cabbage, Chinese cabbage (tight-heading varieties only).
Caneberry	<i>Rubus</i> spp. (including blackberry); <i>Rubus caesius</i> (youngberry); <i>Rubus loganbaccus</i> (loganberry); <i>Rubus idaeus</i> (red and black raspberry); cultivars, varieties, and/or hybrids of these.
Celery	Celery, Florence fennel (sweet anise, sweet fennel, finocchio) (fresh leaves and stalks only).
Cherry	Cherry, sweet, and cherry, tart.
Endive	Endive, escarole.

Fern, edible, fiddlehead	Fern, edible, fiddlehead including: Black lady fern, <i>Deparia japonica</i> (Thunb.) M. Kato; Bracken fern, <i>Pteridium aquilinum</i> (L.) Kuhn; Broad buckler fern, <i>Dryopteris dilatata</i> (Hoffm.) A. Gray; Cinnamon fern, <i>Osmundastrum cinnamomeum</i> (L.) C. Presl; Lady fern, <i>Athyrium filix-femina</i> (L.) Roth ex Mert.; Leather fern, <i>Acrostichum aureum</i> L.; Mother fern, <i>Diplazium proliferum</i> (Lam.) Thouars; Ostrich fern, <i>Matteuccia struthiopteris</i> (L.) Tod.; Vegetable fern, <i>Diplazium esculentum</i> (Retz.) Sw.; Zenmai fern, <i>Osmunda japonica</i> Thunb.
Flowers, edible, multiple species	Acacia blossoms (<i>Acacia senegal</i> (L.) Willd.); Alyssum, sweet (<i>Lobularia maritima</i> (L.) Desv.); Anchusa, garden (<i>Anchusa azurea</i> Mill.); Angelica (<i>Angelica archangelica</i> L.); Apricot, Japanese (<i>Prunus mume</i> Siebold & Zucc.); Arnica (<i>Arnica montana</i> L.); Arugula (<i>Eruca sativa</i> Mill.); Balm (<i>Melissa officinalis</i> L.); Banana (<i>Musa</i> spp.); Basil (<i>Ocimum</i> spp.); Begonia, tuberous (<i>Begonia x tuberhybrida</i> Voss); Bilimbi (<i>Averrhoa bilimbi</i> L.); Bisnaga (<i>Ammi visnaga</i> (L.) Lam.); Blue thistle (<i>Centaurea benedicta</i> (L.) L.); Borage (<i>Borago officinalis</i> L.); Broccoli (<i>Brassica oleracea</i> L. var. <i>italica</i> Plenck); Bugelweed (<i>Lycopus</i> spp.); Burnet (<i>Sanguisorba</i> spp.); Calendula (<i>Calendula officinalis</i> L.); Canadian goldenrod (<i>Solidago canadensis</i> L.); Caper (<i>Capparis spinosa</i> L.); Carambola (<i>Averrhoa carambola</i> L.); Carnation (<i>Dianthus caryophyllus</i> L.); Celandine, greater (<i>Chelidonium majus</i> L.); Chamomile (<i>Chamaemelum</i> spp. and <i>Matricaria</i> spp.); Chaparral (<i>Larrea tridentata</i> (DC.) Coville); Chervil (<i>Anthriscus cerefolium</i> (L.) Hoffm.); Chicory (<i>Cichorium intybus</i> L.); Chive, Chinese (<i>Allium tuberosum</i> Rottler ex Spreng.); Chrysanthemum (<i>Chrysanthemum</i> spp.); Clary (<i>Salvia sclarea</i> L.); Cleavers (<i>Galium aparine</i> L.); Clove (<i>Syzygium aromaticum</i> (L.) Merr. & L. M. Perry); Clover, red (<i>Trifolium pratense</i> L.); Coriander/Cilantro (<i>Coriandrum sativum</i> L.); Cornflower (<i>Centaurea cyanus</i> L.); Costmary (<i>Tanacetum balsamita</i> L. subsp. <i>balsamita</i>); Daisy, English (<i>Bellis perennis</i> L.); Dames rocket (<i>Hesperis matronalis</i> L.); Damiana (<i>Turnera diffusa</i> Willd); Dandelion (<i>Taraxacum officinale</i> F. H. Wigg. aggr.); Daylily (<i>Hemerocallis fulva</i> (L.) L.); Dill (<i>Anethum graveolens</i> L.); Elder (<i>Sambucus nigra</i> L.); Eyebright (<i>Euphrasia</i> spp.); Feijoa (<i>Acca sellowiana</i> (O. Berg) Burret); Fennel (common) (<i>Foeniculum vulgare</i> Mill. subsp. <i>vulgare</i> var. <i>vulgare</i>); Frangipani (<i>Plumeria rubra</i> L.); Fuchsia (<i>Fuchsia</i> spp.); Gardenia (<i>Gardenia jasminoides</i> J. Ellis); Geranium (<i>Pelargonium</i> spp.); Geranium, lemon (<i>Pelargonium crispum</i> (P.J. Bergius) L'Her.); Geranium, rose (<i>Pelargonium graveolens</i> L'Hér.); Ginger, white (<i>Hedychium coronarium</i> J. Koenig); Gladiolus (<i>Gladiolus</i> spp.); Greater periwinkle (<i>Vinca major</i> L.); Hawthorn (<i>Crataegus monogyna</i> Jacq. <i>Crataegus</i> spp.); Hibiscus (<i>Hibiscus</i> spp.); Hibiscus, Chinese (<i>Hibiscus rosa-sinensis</i> L.); Hollyhock (<i>Alcea rosea</i> L.); Honeysuckle, Japanese (<i>Lonicera japonica</i> Thunb.); Horehound (<i>Marrubium vulgare</i> L.); Hyssop (<i>Hyssopus officinalis</i> L.); Hyssop, anise (<i>Agastache foeniculum</i> (Pursh) Kuntze); Impatiens (<i>Impatiens walleriana</i> Hook. f.); Jasmine, Arabian (<i>Jasminum sambac</i> (L.) Aiton); Kewra (<i>Pandanus fascicularis</i> Lam.); Lavender (<i>Lavandula angustifolia</i> Mill.); Lemon (<i>Citrus limon</i> (L.) Burm. f.); Lilac (<i>Syringa vulgaris</i> L.); Lily (<i>Lilium</i> spp.); Lily, mariposa (<i>Calochortus gunnisonii</i> S. Watson); Lily, sego (<i>Calochortus nuttallii</i> Torr. & A. Gray); Lotus (<i>Nelumbo nucifera</i> Gaertn.); Lovage (<i>Levisticum officinale</i> W. D. J. Koch); Mallow, high (<i>Malva sylvestris</i> L.); Marigold (<i>Tagetes</i> spp.); Marjoram (<i>Origanum</i> spp.); Meadowsweet (<i>Filipendula ulmaria</i> (L.) Maxim.); Mint (<i>Mentha</i> spp.); Mioga (<i>Zingiber mioga</i> (Thunb.) Roscoe); Monarda (<i>Monarda</i> spp.); Motherwort (<i>Leonurus cardiaca</i> L.); Mullein (<i>Verbascum thapsus</i> L., <i>Verbascum</i> spp.); Mustard (<i>Brassica</i> spp. and <i>Sinapis</i> spp.); Nasturtium (<i>Tropaeolum</i> spp.); Okra (<i>Abelmoschus esculentus</i> (L.) Moench); Orange, bitter (<i>Citrus aurantium</i> L.); Passion flower (<i>Passiflora</i> spp.); Pea blossoms (<i>Pisum sativum</i> L. subsp. <i>sativum</i> var. <i>sativum</i>); Peach (<i>Prunus persica</i> (L.) Batsch var. <i>persica</i>); Peony, common (<i>Paeonia officinalis</i> L.); Perilla (<i>Perilla frutescens</i> (L.) Britton); Petunia (<i>Petunia x hybrida</i> hort. ex E. Vilm.); Primrose (<i>Primula vulgaris</i> Huds.); Puget sound gumweed (<i>Grindelia integrifolia</i> DC.); Purslane, winter (<i>Claytonia perfoliata</i> Donn ex Willd.); Radish (<i>Raphanus sativus</i> L.); Redbud (<i>Cercis canadensis</i> L.); Rose (<i>Rosa</i> spp.); Rosemary (<i>Rosmarinus officinalis</i> L.); Rose-of-Sharon (<i>Hibiscus syriacus</i> L.); Runner bean, scarlet (<i>Phaseolus coccineus</i> L.); Safflower (<i>Carthamus tinctorius</i> L.); Sage (<i>Salvia officinalis</i> L.); Sage, pineapple (<i>Salvia elegans</i>); Savory, summer (<i>Satureja hortensis</i> L.); Saxifrage, burnet (<i>Pimpinella saxifraga</i> L.); Scotch broom (<i>Cytisus scoparius</i> (L.) Link.); Shepherd's purse (<i>Capsella bursa-pastoris</i> (L.) Medik.); Snapdragon (<i>Antirrhinum majus</i> L.); Sorrel, garden (<i>Rumex acetosa</i> L.); Sorrel, wood (<i>Oxalis acetosella</i> L.); Spilanthes (<i>Blainvillea acmella</i> (L.) Philipson); Squash (<i>Cucurbita</i> spp.); Stock, gillyflower (<i>Matthiola incana</i> (L.) W. T. Aiton); Stoneroot (<i>Collinsonia canadensis</i> L.); Sunflower (<i>Helianthus annuus</i> L.); Sweet william (<i>Dianthus barbatus</i> L.); Sweet wormwood (<i>Artemisia annua</i> L.); Thyme (<i>Thymus vulgaris</i> L.); Tuberose (<i>Polyanthes tuberosa</i> L.); Tulip (<i>Tulipa</i> spp.); Verbena, blue (<i>Verbena hastata</i> L.); Verbena, lemon (<i>Aloysia citrodora</i> Palau); Violet (<i>Viola</i> spp.); Wormwood (<i>Artemisia absinthium</i> L.); Yarrow (<i>Achillea millefolium</i> L.); Yucca (<i>Yucca</i> spp.); and other edible flowers.

Flowers, edible, multiple species	<p>Acacia blossoms (<i>Acacia senegal</i> (L.) Willd.); Alyssum, sweet (<i>Lobularia maritima</i> (L.) Desv.); Anchusa, garden (<i>Anchusa azurea</i> Mill.); Angelica (<i>Angelica archangelica</i> L.); Apricot, Japanese (<i>Prunus mume</i> Siebold & Zucc.); Arnica (<i>Arnica montana</i> L.); Arugula (<i>Eruca sativa</i> Mill.); Balm (<i>Melissa officinalis</i> L.); Banana (<i>Musa</i> spp.); Basil (<i>Ocimum</i> spp.); Begonia, tuberous (<i>Begonia x tuberhybrida</i> Voss); Bilimbi (<i>Averrhoa bilimbi</i> L.); Bisnaga (<i>Ammi visnaga</i> (L.) Lam.); Blue thistle (<i>Centaurea benedicta</i> (L.) L.); Borage (<i>Borago officinalis</i> L.); Broccoli (<i>Brassica oleracea</i> L. var. <i>italica</i> Plenck); Bugelweed (<i>Lycopus</i> spp.); Burnet (<i>Sanguisorba</i> spp.); Calendula (<i>Calendula officinalis</i> L.); Canadian goldenrod (<i>Solidago canadensis</i> L.); Caper (<i>Capparis spinosa</i> L.); Carambola (<i>Averrhoa carambola</i> L.); Carnation (<i>Dianthus caryophyllus</i> L.); Celandine, greater (<i>Chelidonium majus</i> L.); Chamomile (<i>Chamaemelum</i> spp. and <i>Matricaria</i> spp.); Chaparral (<i>Larrea tridentata</i> (DC.) Coville); Chervil (<i>Anthriscus cerefolium</i> (L.) Hoffm.); Chicory (<i>Cichorium intybus</i> L.); Chive, Chinese (<i>Allium tuberosum</i> Rottler ex Spreng.); Chrysanthemum (<i>Chrysanthemum</i> spp.); Clary (<i>Salvia sclarea</i> L.); Cleavers (<i>Galium aparine</i> L.); Clove (<i>Syzygium aromaticum</i> (L.) Merr. & L. M. Perry); Clover, red (<i>Trifolium pratense</i> L.); Coriander/Cilantro (<i>Coriandrum sativum</i> L.); Cornflower (<i>Centaurea cyanus</i> L.); Costmary (<i>Tanacetum balsamita</i> L. subsp. <i>balsamita</i>); Daisy, English (<i>Bellis perennis</i> L.); Dames rocket (<i>Hesperis matronalis</i> L.); Damiana (<i>Turnera diffusa</i> Willd); Dandelion (<i>Taraxacum officinale</i> F. H. Wigg. aggr.); Daylily (<i>Hemerocallis fulva</i> (L.) L.); Dill (<i>Anethum graveolens</i> L.); Elder (<i>Sambucus nigra</i> L.); Eyebright (<i>Euphrasia</i> spp.); Feijoa (<i>Acca sellowiana</i> (O. Berg) Burret); Fennel (common) (<i>Foeniculum vulgare</i> Mill. subsp. <i>vulgare</i> var. <i>vulgare</i>); Frangipani (<i>Plumeria rubra</i> L.); Fuchsia (<i>Fuchsia</i> spp.); Gardenia (<i>Gardenia jasminoides</i> J. Ellis); Geranium (<i>Pelargonium</i> spp.); Geranium, lemon (<i>Pelargonium crispum</i> (P.J. Bergius) L'Her.); Geranium, rose (<i>Pelargonium graveolens</i> L'Hér.); Ginger, white (<i>Hedychium coronarium</i> J. Koenig); Gladiolus (<i>Gladiolus</i> spp.); Greater periwinkle (<i>Vinca major</i> L.); Hawthorn (<i>Crataegus monogyna</i> Jacq. <i>Crataegus</i> spp.); Hibiscus (<i>Hibiscus</i> spp.); Hibiscus, Chinese (<i>Hibiscus rosa-sinensis</i> L.); Hollyhock (<i>Alcea rosea</i> L.); Honeysuckle, Japanese (<i>Lonicera japonica</i> Thunb.); Horehound (<i>Marrubium vulgare</i> L.); Hyssop (<i>Hyssopus officinalis</i> L.); Hyssop, anise (<i>Agastache foeniculum</i> (Pursh) Kuntze); Impatiens (<i>Impatiens walleriana</i> Hook. f.); Jasmine, Arabian (<i>Jasminum sambac</i> (L.) Aiton); Kewra (<i>Pandanus fascicularis</i> Lam.); Lavender (<i>Lavandula angustifolia</i> Mill.); Lemon (<i>Citrus limon</i> (L.) Burm. f.); Lilac (<i>Syringa vulgaris</i> L.); Lily (<i>Lilium</i> spp.); Lily, mariposa (<i>Calochortus gunnisonii</i> S. Watson); Lily, sego (<i>Calochortus nuttallii</i> Torr. & A. Gray); Lotus (<i>Nelumbo nucifera</i> Gaertn.); Lovage (<i>Levisticum officinale</i> W. D. J. Koch); Mallow, high (<i>Malva sylvestris</i> L.); Marigold (<i>Tagetes</i> spp.); Marjoram (<i>Origanum</i> spp.); Meadowsweet (<i>Filipendula ulmaria</i> (L.) Maxim.); Mint (<i>Mentha</i> spp.); Mioga (<i>Zingiber mioga</i> (Thunb.) Roscoe); Monarda (<i>Monarda</i> spp.); Motherwort (<i>Leonurus cardiaca</i> L.); Mullein (<i>Verbascum thapsus</i> L., <i>Verbascum</i> spp.); Mustard (<i>Brassica</i> spp. and <i>Sinapis</i> spp.); Nasturtium (<i>Tropaeolum</i> spp.); Okra (<i>Abelmoschus esculentus</i> (L.) Moench); Orange, bitter (<i>Citrus aurantium</i> L.); Passion flower (<i>Passiflora</i> spp.); Pea blossoms (<i>Pisum sativum</i> L. subsp. <i>sativum</i> var. <i>sativum</i>); Peach (<i>Prunus persica</i> (L.) Batsch var. <i>persica</i>); Peony, common (<i>Paeonia officinalis</i> L.); Perilla (<i>Perilla frutescens</i> (L.) Britton); Petunia (<i>Petunia x hybrida</i> hort. ex E. Vilm.); Primrose (<i>Primula vulgaris</i> Huds.); Puget sound gumweed (<i>Grindelia integrifolia</i> DC.); Purslane, winter (<i>Claytonia perfoliata</i> Donn ex Willd.); Radish (<i>Raphanus sativus</i> L.); Redbud (<i>Cercis canadensis</i> L.); Rose (<i>Rosa</i> spp.); Rosemary (<i>Rosmarinus officinalis</i> L.); Rose-of-Sharon (<i>Hibiscus syriacus</i> L.); Runner bean, scarlet (<i>Phaseolus coccineus</i> L.); Safflower (<i>Carthamus tinctorius</i> L.); Sage (<i>Salvia officinalis</i> L.); Sage, pineapple (<i>Salvia elegans</i>); Savory, summer (<i>Satureja hortensis</i> L.); Saxifrage, burnet (<i>Pimpinella saxifraga</i> L.); Scotch broom (<i>Cytisus scoparius</i> (L.) Link); Shepherd's purse (<i>Capsella bursa-pastoris</i> (L.) Medik.); Snapdragon (<i>Antirrhinum majus</i> L.); Sorrel, garden (<i>Rumex acetosa</i> L.); Sorrel, wood (<i>Oxalis acetosella</i> L.); Spilanthes (<i>Blainvillea acmella</i> (L.) Philipson); Squash (<i>Cucurbita</i> spp.); Stock, gillyflower (<i>Matthiola incana</i> (L.) W. T. Aiton); Stoneroot (<i>Collinsonia canadensis</i> L.); Sunflower (<i>Helianthus annuus</i> L.); Sweet william (<i>Dianthus barbatus</i> L.); Sweet wormwood (<i>Artemisia annua</i> L.); Thyme (<i>Thymus vulgaris</i> L.); Tuberose (<i>Polianthes tuberosa</i> L.); Tulip (<i>Tulipa</i> spp.); Verbena, blue (<i>Verbena hastata</i> L.); Verbena, lemon (<i>Aloysia citrodora</i> Palau); Violet (<i>Viola</i> spp.); Wormwood (<i>Artemisia absinthium</i> L.); Yarrow (<i>Achillea millefolium</i> L.); Yucca (<i>Yucca</i> spp.); and other edible flowers.</p>
Fruit, citrus	Grapefruit, lemon, lime, orange, tangelo, tangerine, citrus citron, kumquat, and hybrids of these.
Garlic	Garlic, great headed; garlic, and serpent garlic.
Guava	Guava (<i>Psidium guajava</i> L.); Guava, cattley (<i>Psidium cattleyanum</i> Sabine); Guava, Para (<i>Psidium acutangulum</i> DC.); Guava, purple strawberry (<i>Psidium cattleyanum</i> Sabine var. <i>cattleyanum</i>); Guava, strawberry (<i>Psidium cattleyanum</i> Sabine var. <i>littorale</i> (Raddi) Fosberg); Guava, yellow strawberry (<i>Psidium cattleyanum</i> Sabine var. <i>cattleyanum</i> forma <i>lucidum</i> O. Deg.)

Lettuce	Lettuce, head; and lettuce, leaf
Lettuce, head	Lettuce, head; crisphead varieties only
Lettuce, leaf	Lettuce, leaf; cos (romaine), butterhead varieties
Marjoram (<i>Origanum</i> spp.); Marjoram, pot (<i>Origanum onites</i> L.); Marjoram, sweet (<i>Origanum majorana</i> L.); Oregano (<i>Origanum vulgare</i> L.)	<i>Origanum</i> spp. (includes sweet or annual marjoram, wild marjoram or oregano, and pot marjoram).
Melon	Muskmelon, including hybrids and/or varieties of <i>Cucumis melo</i> (including true cantaloupe, cantaloupe, casaba, Santa Claus melon, crenshaw melon, honeydew melon, honey balls, Persian melon, golden pershaw melon, mango melon, pineapple melon, snake melon); and watermelon, including hybrids and/or varieties of (<i>Citrullus</i> spp.).
Mint (<i>Mentha</i> spp.)	Mint (<i>Mentha</i> spp.); Applemint (<i>Mentha suaveolens</i> Ehrh.); Horsemint (<i>Mentha longifolia</i> (L.) Huds.); Mint, corn (<i>Mentha arvensis</i> L.); Peppermint (<i>Mentha x piperita</i> L.); Spearmint, (<i>Mentha spicata</i> L.); Spearmint, Scotch (<i>Mentha x gracilis</i> Sole); Watermint (<i>Mentha aquatica</i> L.); Pennyroyal (<i>Mentha pulegium</i> L.)
Muskmelon	<i>Cucumis melo</i> (includes true cantaloupe, cantaloupe, casaba, Santa Claus melon, crenshaw melon, honeydew melon, honey balls, Persian melon, golden pershaw melon, mango melon, pineapple melon, snake melon, and other varieties and/or hybrids of these.)
Onion	Bulb onion; green onion; and garlic.
Onion, bulb	Bulb onion; garlic; great headed garlic; serpent garlic; Chinese onion; pearl onion; potato onion; and shallot, bulb.
Onion, green	Green onion; lady's leek; leek; wild leek; Beltsville bunching onion; fresh onion; tree onion, tops; Welsh onion; and shallot, fresh leaves.
Palm hearts	Palm hearts, various species, including: African fan palm, <i>Borassus aethiopum</i> Mart.; Cabbage palm, <i>Euterpe oleracea</i> Mart.; Cabbage palmetto, <i>Sabal palmetto</i> (Walter) Schult. & Schult. f.; Coconut, <i>Cocos nucifera</i> L.; Palmyra palm, <i>Borassus flabellifera</i> L.; Peach Palm, <i>Bactris gasipaes</i> Kunth; Royal palm, <i>Roystonea oleracea</i> (Jacq.) O.F. Cook; Salak palm, <i>Salacca zalacca</i> (Gaertn.) Voss; Saw palmetto, <i>Serenoa repens</i> (W. Bartram) Small; Wine palm, <i>Raphia</i> spp.
Peach	Peach, nectarine

Pea	<i>Cajanus cajan</i> (pigeon pea); <i>Cicer arietinum</i> (chickpea, garbanzo bean); <i>Lens culinaris</i> (lentil); Grass pea; <i>Pisum</i> spp. (including, but not limited to dry pea, dwarf pea, English pea, field pea, garden pea, green pea, marrowfat pea, snap pea, snow pea, sugar snap pea, wrinkled pea and yellow pea); cultivars, varieties and/or hybrids of these commodities. [Note: A variety of pesticide tolerances have been previously established for pea and/or bean. Chickpea/garbanzo bean is also listed in the definition for “bean”. For garbanzo bean/chickpea only, the highest established pea or bean tolerance will apply to pesticide residues found in this commodity]. <i>Lens culinaris</i> (lentil); <i>Pisum</i> spp. (includes dwarf pea, garden pea, green pea, English pea, field pea, and edible pod pea). [Note: A variety of pesticide tolerances have been previously established for pea and/or bean. Chickpea/garbanzo bean is now classified in both the bean and the pea categories. For garbanzo bean/chickpea only, the highest established pea or bean tolerance will apply to pesticide residues found in this commodity.]
Pea, dry, seed	All peas in the entry “Pea” in dry seed form.
Pea, edible podded	All peas in the entry “Pea” in edible podded form.
Pea, succulent	All peas in the entry “Pea” in edible podded or succulent shelled form.
Pea, succulent shelled	All peas in the entry “Pea” in succulent shelled form.
Pepper	All varieties of pepper including pimento and bell, hot, and sweet pepper.
Radish, oriental, roots	<i>Raphanus sativus</i> var. <i>longipinnatus</i> (roots and tops), including Chinese or Japanese radish (both white and red), winter radish, daikon, lobok, lo pak, and other cultivars and/or hybrids of these.
Radish, oriental, tops)	<i>Raphanus sativus</i> var. <i>longipinnatus</i> (roots and tops), including Chinese or Japanese radish (both white and red), winter radish, daikon, lobok, lo pak, and other cultivars and/or hybrids of these.
Rapeseed	<i>Brassica napus</i> , <i>B. campestris</i> , and <i>Crambe abyssinica</i> (oilseed-producing varieties only which include canola and crambe.)
Raspberry	<i>Rubus</i> spp. (including bababerry; black raspberry; blackcap; caneberry; framboise; frambueso; himbeere; keriberry; mayberry; red raspberry; thimbleberry; tulameen; yellow raspberry; and cultivars, varieties, and/or hybrids of these).
Sorghum, grain, grain	<i>Sorghum</i> spp. [sorghum, grain, sudangrass (seed crop), and hybrids of these grown for its seed].
Sorghum, forage, stover	<i>Sorghum</i> spp. [sorghum, forage; sorghum, stover; sudangrass, and hybrids of these grown for forage and/or stover.
Squash	Pumpkin, summer squash, and winter squash.
Sugar apple	<i>Annona squamosa</i> L. and its hybrid atemoya (<i>Annona cherimola</i> Mill X <i>A. squamosa</i> L.) Also includes true custard apple (<i>Annona reticulata</i> L.).

Squash, summer	Fruits of the gourd (<i>Cucurbitaceae</i>) family that are consumed when immature, 100% of the fruit is edible either cooked or raw, once picked it cannot be stored, has a soft rind which is easily penetrated, and if seeds were harvested they would not germinate; e.g., <i>Cucurbita pepo</i> (i.e., crookneck squash, straightneck squash, scallop squash, and vegetable marrow); <i>Lagenaria</i> spp. (i.e., spaghetti squash, hyotan, cucuzza); <i>Luffa</i> spp. (i.e., hechima, Chinese okra); <i>Momordica</i> spp. (i.e., bitter melon, balsam pear, balsam apple, Chinese cucumber); <i>Sechium edule</i> (chayote); and other cultivars and/or hybrids of these.
Sweet potato	Sweet potato, yam.
Tangerine	Tangerine (mandarin or mandarin orange); clementine; Mediterranean mandarin; satsuma mandarin; tangelo; tangor; cultivars, varieties, and/or hybrids of these.
Tomato	Tomato, tomatillo.
Turnip tops or turnip greens	Broccoli raab (raab, raab salad), hanover salad, turnip tops (turnip greens).
Wheat	Wheat, triticale.

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)