

40 C.F.R. § 180.317

Propyzamide; tolerances for residues.

(a) *General.* Tolerances are established for residues of the herbicide propyzamide, including its metabolites and degradates, in or on the commodities in the table in this paragraph. Compliance with the tolerance levels specified in this paragraph is to be determined by measuring only those propyzamide residues convertible to methyl 3,5-dichlorobenzoate, expressed as the stoichiometric equivalent of propyzamide, 3,5-dichloro-*N*-(1,1-dimethyl-2-propynyl)benzamide, in or on the commodity.

Commodity	Parts per million
Alfalfa, seed	10.0
Animal feed, nongrass, group 18	10.0
Apple	0.1
Artichoke, globe	0.01
Blackberry	0.05
Blueberry	0.05
Boysenberry	0.05
Cattle, fat	0.2
Cattle, kidney	0.4
Cattle, liver	0.4
Cattle, meat	0.02
Cattle, meat byproducts, except kidney and liver	0.02
Egg	0.02
Endive	1.0
Fruit, stone, group 12	0.1
Goat, fat	0.2
Goat, kidney	0.4

Goat, liver	0.4
Goat, meat	0.02
Goat, meat byproducts, except kidney and liver	0.02
Grape	0.1
Hog, fat	0.2
Hog, kidney	0.4
Hog, liver	0.4
Hog, meat	0.02
Hog, meat byproducts, except kidney and liver	0.02
Horse, fat	0.2
Horse, kidney	0.4
Horse, liver	0.4
Horse, meat	0.02
Horse, meat byproducts, except kidney and liver	0.02
Lettuce, head	1.0
Lettuce, leaf	1.0
Milk	0.02
Pear	0.1
Poultry, fat	0.02
Poultry, liver	0.2
Poultry, meat	0.02
Poultry, meat byproducts, except liver	0.02
Radicchio	2.0
Raspberry	0.05
Sheep, fat	0.2
Sheep, kidney	0.4
Sheep, liver	0.4
Sheep, meat	0.02

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)