

40 C.F.R. § 180.399

Iprodione; tolerances for residues.

(a) *General.* (1) Tolerances are established for the combined residues of the fungicide iprodione [3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide], its isomer 3-(1-methylethyl)-N-(3,5-dichlorophenyl)-2,4-dioxo-1-imidazolidinecarboxamide, and its metabolite 3-(3,5-dichlorophenyl)-2,4-dioxo-1-imidazolidine-carboxamide in or on the following food commodities:

Commodity	Parts per million
Almond, hulls	2.0
Almond	0.3
Apricot	20.0
Bean, dry, seed	2.0
Bean, forage	90.0
Bean, succulent	2.0
Blueberry	15.0
Boysenberry	15.0
Broccoli	25.0
Caneberry subgroup 13A	25.0
Carrot, roots	5.0
Cherry, sweet, postharvest	20.0
Cherry, tart	20.0
Cotton, undelinted seed	0.10
Cowpea, hay	90.0
Currant	15.0
Garlic	0.1
Ginseng	2.0

Ginseng, dried root	4.0
Grape	60.0
Grape, raisin	300
Kiwifruit	10.0
Lettuce	25.0
Nectarine, postharvest	20.0
Onion, bulb	0.5
Peach, postharvest	20.0
Peanut	0.5
Peanut, hay	150.0
Plum, postharvest	20.0
Plum, prune	20.0
Potato	0.5
Raspberry	15.0
Rice, bran	30.0
Rice, grain	10.0
Rice, hulls	50.0
Strawberry	15.0

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)