

# 40 C.F.R. § 180.430

## Fenoxaprop-ethyl; tolerances for residues.

(a) *General.* Tolerances are established for residues of the herbicide fenoxaprop-ethyl, including its metabolites and degradates, in or on the commodities in the table in this paragraph. Compliance with the tolerance levels specified in this paragraph is to be determined by measuring only the sum of fenoxaprop-ethyl, ( $\pm$ )-ethyl 2-[4-[(6-chloro-2-benzoxazolyl)oxy]phenoxy]propanoate, and its metabolites, 2-[4-[(6-chloro-2-benzoxazolyl)oxy]phenoxy]propanoic acid and 6-chloro-2,3-dihydrobenzoxazol-2-one, calculated as the stoichiometric equivalent of fenoxaprop-ethyl, in or on the commodity.

Commodity	Parts per million
Barley, grain	0.05
Barley, straw	0.1
Cattle, fat	0.05
Cattle, meat byproducts	0.05
Cattle, meat	0.05
Cotton, undelinted seed	0.05
Goat, fat	0.05
Goat, meat byproducts	0.05
Goat, meat	0.05
Hog, fat	0.05
Hog, meat byproducts	0.05
Hog, meat	0.05
Horse, fat	0.05
Horse, meat byproducts	0.05
Horse, meat	0.05
Milk	0.02
Peanut	0.05

Peanut, hulls	0.05
Rice, grain	0.05
Sheep, fat	0.05
Sheep, meat byproducts	0.05
Sheep, meat	0.05
Soybean	0.05
Wheat, grain	0.05
Wheat, straw	0.50

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)