

40 C.F.R. § 180.434

Propiconazole; tolerances for residues.

(a) *General.* (1) Tolerances are established for residues of propiconazole, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only those propiconazole residues convertible to 2,4-dichlorobenzoic acid (2,4-DCBA), expressed as the stoichiometric equivalent of propiconazole, in or on the commodity in the table below:

Commodity	Parts per million
Almond, hulls	7.0
Avocado	0.2
Banana	0.2
Barley, bran	6.0
Barley, grain	3.0
Barley, hay	30
Barley, straw	20
Bean, dry seed	0.40
Bean, snap	0.70
Bean, succulent shelled	0.10
Beet, garden, tops	5.5
Beet, sugar, dried pulp	1.0
Beet, sugar, molasses	1.5
Beet, sugar, roots	0.3
Beet, sugar, tops	10
Brassica, leafy greens, subgroup 4-16B, except watercress	20
Bushberry, subgroup 13-07B	1.0

Caneberry, subgroup 13-07A	1.0
Cattle, fat	0.05
Cattle, kidney	2.0
Cattle, liver	2.0
Cattle, meat	0.05
Cattle, meat byproducts, except liver and kidney	0.05
Celtuce	5
Cilantro, leaves	13
Citrus, oil	1000
Corn, field, forage	12
Corn, field, grain	0.2
Corn, field, stover	30
Corn, pop, grain	0.2
Corn, pop, stover	30
Corn, sweet, forage	6.0
Corn, sweet, kernel plus cob with husks removed	0.1
Corn, sweet, stover	30
Dill, seed	15
Dillweed, dried leaves	80
Dillweed, fresh leaves	30
Fennel, Florence, fresh leaves and stalk	5
Fruit, citrus, group 10-10	8.0
Fruit, stone, group 12-12, except plum	4.0
Goat, fat	0.05
Goat, kidney	2.0
Goat, liver	2.0
Goat, meat	0.05
Goat, meat byproducts, except liver and kidney	0.05

Grain, aspirated fractions	110
Grass, forage	0.5
Grass, hay	0.5
Grass, straw	40
Hog, kidney	0.2
Hog, liver	0.2
Horse, fat	0.05
Horse, kidney	2.0
Horse, liver	2.0
Horse, meat	0.05
Horse, meat byproducts, except liver and kidney	0.05
Leaf petiole vegetable subgroup 22B	5
Low growing berry subgroup 13-07G, except cranberry	1.3
Milk	0.05
Mushroom	0.1
Nut, tree, group 14-12	0.10
Oat, forage	4.0
Oat, grain	3.0
Oat, hay	15
Oat, straw	10
Onion, bulb subgroup 3-07A	0.2
Onion, green, subgroup 3-07B	9.0
Parsley, fresh leaves	13
Parsley, dried leaves	35
Peanut	0.2
Peanut, hay	20
Peppermint, tops	10.0
Pineapple	4.5

Pineapple, process residue	7.0
Plum	0.60
Rapeseed subgroup 20A	0.30
Quinoa, grain	3.0
Radish, tops	0.20
Rice, bran	15
Rice, grain	7.0
Rice, hulls	20
Rye, bran	0.6
Rye, forage	9.0
Rye, grain	0.3
Rye, straw	10
Sheep, fat	0.05
Sheep, kidney	2.0
Sheep, liver	2.0
Sheep, meat	0.05
Sheep, meat byproducts, except liver and kidney	0.05
Sorghum, grain, forage	12
Sorghum, grain, grain	3.5
Sorghum, grain, stover	15
Soybean, forage	11
Soybean, hay	30
Soybean, seed	2.0
Spearmint, tops	10.0
Sugarcane, cane	0.4
Swiss chard	5
Ti palm, leaves	10
Ti palm, roots	0.30

Tomato	3.0
Vegetable, foliage of legume, group 7	30
Vegetable, root, except sugar beet, subgroup 1B	0.3
Watercress	6.0
Wheat, bran	0.6
Wheat, forage	15
Wheat, grain	0.3
Wheat, hay	30
Wheat, straw	20

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)