

40 C.F.R. § 180.448

Hexythiazox; tolerance for residues.

(a) *General.* Tolerances are established for residues of hexythiazox, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only hexythiazox and its metabolites containing the (4-chlorophenyl)-4-methyl-2-oxo-3-thiazolidine moiety, calculated as the stoichiometric equivalent of hexythiazox.

Commodity	Parts per million
Almond, hulls	10
Apple, wet pomace	0.40
Beet, sugar, dried pulp	0.30
Beet, sugar, root	0.15
Berry, low growing, subgroup 13-07G	6
Caneberry, Subgroup 13-07A	3
Cattle, fat	0.05
Cattle, meat byproducts	0.5
Citrus, oil	25
Corn, field, forage	3.0
Corn, field, grain	0.02
Corn, field, stover	7.0
Cotton, gin byproducts	15
Cotton, undelinted seed	0.4
Date, dried	3
Egg	0.05
Fruit, pome, group 11-10	0.4
Fruit, small, vine climbing, subgroup 13-07F, except fuzzy kiwifruit	1

Fruit, stone, group 12	1.0
Goat, fat	0.05
Goat, meat byproducts	0.5
Grain, aspirated fractions	5
Hog, fat	0.02
Hog, meat byproducts	0.05
Hop, dried cones	20
Horse, fat	0.05
Horse, meat byproducts	0.5
Milk	0.05
Nut, tree, group 14	0.30
Pepper/eggplant subgroup 8-10B	1.5
Peppermint, tops	2.0
Pistachio	0.30
Plum, prune, dried	1.3
Poultry, fat	0.05
Poultry, meat byproducts	0.05
Sheep, fat	0.05
Sheep, meat byproducts	0.5
Spearmint, tops	2.0
Tea, dried ¹	15
Tomato	0.50

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)