

40 C.F.R. § 180.463

Quinclorac; tolerances for residues.

(a)

(1) *General.* Tolerances are established for residues of the herbicide quinclorac, including its metabolites and degradates, in or on the commodities in the table in this paragraph. Compliance with the tolerance levels specified in this paragraph is to be determined by measuring only quinclorac, 3,7-dichloro-8-quinolinecarboxylic acid, in or on the commodity.

Table 1 to Paragraph (a)(1)

Commodity	Parts per million
Asparagus	0.08
Bushberry, subgroup 13-07B	0.08
Caneberry subgroup 13-07A	0.08
Barley, grain	2.0
Berry, low growing, except strawberry, subgroup 13-07H	1.5
Cattle, fat	0.7
Cattle, meat byproducts	1.5
Cattle, meat	0.05
Egg	0.05
Goat, fat	0.7
Goat, meat byproducts	1.5
Goat, meat	0.05
Grain, aspirated fractions	1200
Grass, forage	150
Grass, hay	130
Hog, fat	0.7

Hog, meat byproducts	1.5
Hog, meat	0.05
Horse, fat	0.7
Horse, meat byproducts	1.5
Horse, meat	0.05
Milk	0.05
Poultry, fat	0.05
Poultry, meat byproducts	0.1
Poultry, meat	0.05
Rhubarb	0.5
Rice, bran	30
Rice, grain	10
Sheep, fat	0.7
Sheep, meat byproducts	1.5
Sheep, meat	0.05
Sorghum, grain, forage	3.0
Sorghum, grain, grain	6.0
Sorghum, grain, stover	1.0
Wheat, forage	1.0
Wheat, germ	0.75
Wheat, grain	0.5
Wheat, hay	0.5
Wheat, straw	0.1

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)