

40 C.F.R. § 180.478

Rimsulfuron; tolerances for residues.

(a) *General.* Tolerances are established for residues of the herbicide rimsulfuron, including its metabolites and degradates, in or on the commodities in the following table. Compliance with the tolerance levels specified in the following table is to be determined by measuring only rimsulfuron, N-[[[4,6-dimethoxy-2-pyrimidinyl)amino] carbonyl]-3-(ethylsulfonyl)-2-pyridinesulfonamide.

Commodity	Parts per million
Almond, hulls	0.09
Berry, low growing, except strawberry, subgroup 13-07H	0.02
Bushberry, subgroup 13-07B	0.01
Caneberry, subgroup 13-07A	0.01
Chicory, roots	0.01
Chicory, tops	0.01
Corn, field, forage	0.4
Corn, field, grain	0.1
Corn, field, stover	2.5
Fruit, citrus, group 10-10	0.01
Fruit, pome, group 11-10	0.01
Fruit, small, vine climbing, except fuzzy kiwifruit, subgroup 13-07F	0.01
Fruit, stone, group 12-12	0.01
Grain, aspirated fractions	4.5
Nut, tree, group 14-12	0.01
Pomegranate	0.01
Sorghum, grain, forage	0.01
Sorghum, grain, grain	0.01

Sorghum, grain, stover	0.01
Soybean, forage	0.25
Soybean, hay	1.2
Soybean, hulls	0.04
Soybean, seed	0.01
Tomato	0.05
Tropical and subtropical, small fruit, edible peel, subgroup 23A	0.01
Vegetable, tuberous and corm, subgroup 1C	0.10

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)