

40 C.F.R. § 180.484

Flutolanil; tolerances for residues.

(a) *General.* Tolerances are established for residues of flutolanil, *N*-(3-(1-methylethoxy) phenyl)-2-(trifluoromethyl)benzamide, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only flutolanil and its metabolites converted to 2-(trifluoromethyl) benzoic acid and calculated as flutolanil, in or on the following commodities:

Commodity	Parts per million
Cattle, fat	0.10
Cattle, kidney	1.00
Cattle, liver	2.00
Cattle, meat byproducts	0.05
Cattle, meat	0.05
Cotton, gin byproducts	0.20
Cotton, undelinted seed	0.20
Egg	0.05
Goat, fat	0.10
Goat, kidney	1.00
Goat, liver	2.00
Goat, meat byproducts	0.05
Goat, meat	0.05
Hog, fat	0.10
Hog, kidney	1.00
Hog, liver	2.00
Hog, meat byproducts	0.05

Hog, meat	0.05
Horse, fat	0.10
Horse, kidney	1.00
Horse, liver	2.00
Horse, meat byproducts	0.05
Horse, meat	0.05
Milk	0.05
Peanut	0.5
Peanut, hay	15.0
Peanut, meal	1.0
Potato	0.20
Potato, wet peel	0.30
Poultry, fat	0.05
Poultry, meat	0.05
Poultry, meat byproducts	0.05
Rice, bran	10.0
Rice, grain	7.0
Rice, hulls	25.0
Sheep, fat	0.10
Sheep, kidney	1.00
Sheep, liver	2.00
Sheep, meat	0.05
Sheep, meat byproducts	0.05
Soybean, forage	8.0
Soybean, hay	2.5
Soybean, seed	0.20
Turnip, greens	0.1
Vegetable, brassica, leafy, group 5	0.1

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)