

# 40 C.F.R. § 180.523

## Metaldehyde; tolerances for residues.

(a) *General.* Tolerances are established for residues of the molluscicide metaldehyde, including its metabolites and degradates, in or on the commodities listed in the following table. Compliance with the specified tolerance levels is to be determined by measuring only metaldehyde, 2,4,6,8-tetramethyl-1,3,5,7-tetroxocane, in or on the commodity.

Commodity	Parts per million
Artichoke, globe	0.07
Berry, low growing, subgroup 13-07G	6.25
Bushberry subgroup 13-07B	0.15
Cactus	0.07
Caneberry subgroup 13-07A	0.15
Corn, field, forage	0.30
Corn, field, grain	0.05
Corn, field, stover	0.10
Corn, sweet, forage	0.30
Corn, sweet, kernel plus cob with husks removed	0.05
Corn, sweet, stover	0.10
Fruit, citrus, group 10-10	0.26
Ginseng	0.05
Grass, forage	2.0
Grass, hay	2.0
Leaf petioles subgroup 4B	0.50
Lettuce	1.73
Pea and bean, succulent shelled, subgroup 6B	0.20

Peppermint, oil	12
Peppermint, tops	4.0
Spearmint, oil	12
Spearmint, tops	4.0
Taro, corm	0.15
Taro, leaves	1.0
Tomato subgroup 8-10A	0.24
Vegetable, brassica, leafy, group 5	2.5
Vegetable, foliage of legume, except soybean, subgroup 7A	1.5
Vegetable, legume, edible podded subgroup 6A	0.80
Watercress	3.2

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)