

40 C.F.R. § 180.572

Bifenazate; tolerance for residues.

(a) *General.* (1) Tolerances are established for residues of bifenazate (1-methylethyl 2-(4-methoxy[1,1'-biphenyl]-3-yl)hydrazinecarboxylate) including its metabolites and degradates, in or on the commodities listed in the following table. Compliance with the tolerance levels specified are to be determined by measuring only the sum of bifenazate and its metabolite, diazinecarboxylic acid, 2-(4-methoxy-[1,1'-biphenyl]-3-yl), 1-methylethyl ester, (calculated as the stoichiometric equivalent of bifenazate) in or on the following food commodities:

Commodity	Parts per million
Acerola	0.90
Almond, hulls	15
Apple, wet pomace	1.2
Atemoya	1.6
Avocado	7.0
Bean, dry seed	0.60
Berry, low-growing subgroup 13-07G	1.5
Biriba	1.6
Black sapote	7.0
Caneberry subgroup 13-07A	5.0
Canistel	7.0
Cattle, fat	0.10
Cherimoya	1.6
Cotton, gin byproducts	35
Cotton, undelinted seed	0.75
Custard apple	1.6
Feijoa	0.90

Fruit, pome, group 11-10	0.7
Fruit, small, vine climbing subgroup 13-07F, except fuzzy kiwifruit	1.0
Fruit, stone, group 12, except plum	2.5
Goat, fat	0.10
Grape	0.75
Grape, raisin	1.2
Guava	0.9
Herb, subgroup 19A, except chervil and chive	300
Hog, fat	0.10
Hop, dried cones	15
Horse, fat	0.10
Jaboticaba	0.90
Ilama	1.6
Longan	5.0
Lychee	5.0
Mango	7.0
Nut, tree, group 14	0.20
Okra	2.0
Papaya	7.0
Passionfruit	0.90
Pea and bean, succulent shelled, subgroup 6B	0.70
Peppermint, tops	25
Pistachio	0.20
Plum	0.20
Pulasan	5.0
Rambutan	5.0
Sapodilla	7.0
Sapote, mamey	7.0

Sheep, fat	0.10
Soursop	1.6
Soybean, succulent shelled	0.70
Spanish lime	5.0
Spearmint, tops	25
Star apple	7.0
Starfruit	0.90
Strawberry	1.5
Sugar apple	1.6
Vegetable, cucurbit, group 9	0.75
Vegetable, fruiting, group 8-10	4.0
Vegetable, legume, edible-podded, subgroup 6A	6.0
Vegetable, tuberous and corm, subgroup 1C	0.10
Wax jambu	0.90

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)