

40 C.F.R. § 180.573

Tepraloxymid; tolerances for residues.

(a) *General.* (1) Tolerances are established for residues of tepraloxymid, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only the combined residues of tepraloxymid, (2-[1-[[[(2E)-3-chloro-2-propen-1-yl]oxy]imino]propyl]-3-hydroxy-5-(tetrahydro-2H-pyran-4-yl)-2-cyclohexen-1-one) and its metabolites convertible to GP (3-(tetrahydropyran-4-yl)pentane-1,5-dioic acid) and OH-GP (3-hydroxy-3-(tetrahydropyran-4-yl)pentane-1,5-dioic acid), calculated as tepraloxymid, in or on the commodities.

Commodity	Parts per million	Expiration/revocation date
Cotton, undelinted seed	0.2	12/31/18
Cotton, gin byproducts	3.0	12/31/18
Flax, seed	0.10	12/31/18
Grain, aspirated fraction	1200.0	12/31/18
Pea and bean, dried shelled, except soybean, subgroup 6C ₁	0.10	12/31/18
Soybean, seed	6.0	12/31/18
Soybean, hulls	8.0	12/31/18
Sunflower subgroup 20B ₁	0.20	12/31/18

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)