

# 40 C.F.R. § 180.578

## Acetamiprid; tolerances for residues.

(a) *General.* (1) Tolerances are established for residues of the insecticide acetamiprid (1E)-N-[(6-chloro-3-pyridinyl)methyl]-N'-cyano-N-methylethanimidamide, including its metabolites and degradates, in or on the commodities in the table below as a result of the application of acetamiprid. Compliance with the tolerance levels specified below is to be determined by measuring only acetamiprid in or on the following commodities.

Commodity	Parts per million
Almond, hulls	5.0
Asparagus	0.80
Berry, low growing subgroups 13-07G	0.60
Brassica, leafy greens, subgroup 4-16B	15
Bushberry subgroup 13-07B	1.6
Caneberry subgroup 13-07A	1.6
Celtuce	3
Citrus, dried pulp	1.20
Corn, sweet, forage	15
Corn, sweet, kernel plus cob with husks removed	0.01
Corn, sweet, stover	30
Cotton, gin byproducts	20.0
Cottonseed subgroup 20C	0.7
Fennel, florence, fresh leaves and stalk	3
Fruit, citrus, group 10-10	1.0
Fruit, pome, group 11-10	1.0
Fruit, small, vine climbing, except fuzzy kiwifruit, subgroup 13-07F	0.35

Fruit, stone, group 12-12	1.5
Grain, aspirated fractions	5.0
Kohlrabi	1.2
Leaf petiole vegetable subgroup 22B	3
Leafy greens subgroup 4-16A	3
Nut, tree, group 14-12	0.1
Onion, bulb, subgroup 3-07A	0.02
Onion, green, subgroup 3-07B	4.5
Pea and bean, succulent shelled, subgroup 6B	0.40
Rapeseed subgroup 20A	0.01
Soybean, hulls	0.04
Soybean, seed	0.03
Tea, dried <sup>1</sup>	50.0
Tomato, paste	0.40
Tropical and subtropical, medium to large fruit, smooth, inedible peel, subgroup 24B	0.5
Vegetable, <i>brassica</i> , head and stem, group 5-16	1.2
Vegetable, cucurbit, group 9	0.50
Vegetable, fruiting, group 8-10	0.20
Vegetable, legume, edible podded, subgroup 6A	0.60
Vegetable, tuberous and corm, group 1	0.01

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)