

40 C.F.R. § 180.585

Pyraflufen-ethyl; tolerances for residues.

(a) *General*. Tolerances are established for residues of the herbicide, pyraflufen-ethyl, including its metabolites and degradates, in the commodities in the table below. Compliance with the plant commodity tolerance levels specified in the table is to be determined by measuring only the sum of the parent pyraflufen-ethyl, ethyl 2-[2-chloro-5-(4-chloro-5-difluoromethoxy)-1-methyl-1H-pyrazol-3-yl]-4-fluorophenoxy] acetate, and its acid metabolite, E-1, 2-chloro-5-(4-chloro-5-difluoromethoxy-1-methyl-1H-pyrazol-3-yl)-4-fluorophenoxyacetic acid, calculated as the stoichiometric equivalent of pyraflufen-ethyl in or on the commodity. Compliance with the livestock commodity tolerance levels specified in the table is to be determined by measuring only the sum of the parent pyraflufen-ethyl, ethyl 2-[2-chloro-5-(4-chloro-5-difluoromethoxy)-1-methyl-1H-pyrazol-3-yl]-4-fluorophenoxy] acetate and its acid metabolites: E-1, 2-chloro-5-(4-chloro-5-difluoromethoxy-1-methyl-1H-pyrazol-3-yl)-4-fluorophenoxyacetic acid, and E-9, 2-chloro-5-(4-chloro-5-difluoromethoxy-1H-pyrazol-3-yl)-4-fluorophenoxyacetic acid, both calculated as the stoichiometric equivalent of pyraflufen-ethyl in or on the commodity.

This document is only available to subscribers. Please log in or purchase access.

Purchase Login