

40 C.F.R. § 180.593

Etoxazole; tolerances for residues.

(a) *General.* Tolerances are established for residues of etoxazole, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only etoxazole (2-(2,6-difluorophenyl)-4-[4-(1,1-dimethylethyl)-2-ethoxyphenyl]-4,5-dihydrooxazole) in or on the commodity.

Commodity	Parts per million
Almond, hulls	2.0
Apple, wet pomace	0.50
Avocado	0.20
Beet, sugar, leaves	1
Beet, sugar, roots	0.02
Berry, low growing, subgroup 13-07G	0.50
Caneberry subgroup 13-07A	1.5
Canistel	0.20
Cattle, fat	0.02
Cattle, liver	0.01
Cherry subgroup 12-12A	1.0
Corn, field, forage	0.80
Corn, field, grain	0.01
Corn, field, refined oil	0.03
Corn, field, stover	4.0
Corn, pop, grain	0.01
Corn, pop, stover	4.0
Corn, sweet, forage	1.5

Corn, sweet, kernel plus cob with husks removed	0.01
Corn, sweet, stover	5.0
Cotton, gin byproducts	1.0
Cottonseed subgroup 20C	0.05
Fruit, pome, group 11-10	0.20
Fruit, small vine climbing, except fuzzy kiwifruit, subgroup 13-07F	0.50
Goat, fat	0.02
Goat, liver	0.01
Grape, raisin	1.5
Hop, dried cones	7.0
Horse, fat	0.02
Horse, liver	0.01
Mango	0.20
Melon subgroup 9A	0.20
Milk, fat	0.01
Nut, tree group 14-12	0.01
Orange 2	0.10
Orange, oil 2	1.0
Papaya	0.20
Peach subgroup 12-12B	1.0
Pepper/eggplant subgroup 8-10B	0.20
Peppermint, oil	20
Peppermint, tops	10
Plum, prune, dried	0.30
Plum subgroup 12-12C	0.15
Sapodilla	0.20
Sapote, black	0.20
Sapote, mamey	0.20

Sheep, fat	0.02
Sheep, liver	0.01
Soybean, seed	0.02
Spearmint, oil	20
Spearmint, tops	10
Squash/cucumber subgroup 9B	0.02
Star apple	0.20
Tangerine 1	0.10
Tea, dried *	15
Tomato	0.20

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)