

40 C.F.R. § 180.609

Fluoxastrobin; tolerances for residues.

(a) *General.* (1) Tolerances are established for residues of fluoxastrobin, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only fluoxastrobin, (1E)-[2-[[6-(2-chlorophenoxy)-5-fluoro-4-pyrimidinyl]oxy]phenyl](5,6-dihydro-1,4,2-dioxazin-3-yl)methanone O-methyloxime and its Z isomer, (1Z)-[2-[[6-(2-chlorophenoxy)-5-fluoro-4-pyrimidinyl]oxy]phenyl](5,6-dihydro-1,4,2-dioxazin-3-yl)methanone O-methyloxime, calculated as the stoichiometric equivalent of fluoxastrobin.

Commodity	Parts per million
Avocado	1.0
Barley, grain	0.40
Barley, hay	15
Barley, straw	15
Berry, low growing, subgroup 13-07G	1.9
Corn, field, forage	3.0
Corn, field, grain	0.02
Corn, field, stover	4.5
Corn, sweet, forage	13
Corn, sweet, kernel plus cob with husks removed	0.01
Corn, sweet, stover	10
Cotton, gin byproducts	0.01
Cotton, undelinted seed	0.01
Grain, aspirated grain fractions	60
Leaf petioles subgroup 4B	4.0
Melon subgroup 9A	1.5
Pea and bean, dried shelled, except soybean, subgroup 6C	0.20

Peanut	0.02
Peanut, hay	20.0
Peanut, refined oil	0.06
Rapeseed, subgroup 20A	0.70
Rice, grain	4.0
Sorghum, grain, forage	5.0
Sorghum, grain, grain	1.5
Sorghum, grain, stover	5.0
Soybean, forage	9.0
Soybean, hay	1.2
Soybean, hulls	0.20
Soybean, seed	0.05
Squash/cucumber subgroup 9B	0.50
Tomato, paste	1.5
Vegetable, fruiting, group 8	1.0
Vegetable, tuberous and corm, subgroup 1C	0.010
Wheat, forage	7.0
Wheat, grain	0.15
Wheat, hay	17
Wheat, straw	11

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)