

# 40 C.F.R. § 180.613

## Flonicamid; tolerances for residues.

(a) *General.* (1) Tolerances are established for the residues of the insecticide flonicamid, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only the sum of flonicamid, *N*-(cyanomethyl)-4-(trifluoromethyl)-3-pyridinecarboxamide, and its metabolites, TFNA (4-trifluoromethylnicotinic acid), TFNA-AM (4-trifluoromethylnicotinamide), and TFNG, *N*-(4-trifluoromethylnicotinoyl)glycine, calculated as the stoichiometric equivalent of flonicamid, in or on the following commodities.

Table 1 to Paragraph (a)(1)

| Commodity   | Parts per million |
|---|-------------------|
| Alfalfa, forage   | 10.0              |
| Alfalfa, hay  | 7                 |
| Alfalfa, seed   | 1.5               |
| Almond, hulls   | 9.0               |
| Berry, low-growing, subgroup 13-07G                                 | 2                 |
| <i>Brassica</i> , leafy greens, subgroup 4-16B, except radish, tops | 16                |
| Bushberry subgroup 13-07B   | 1.5               |
| Caneberry subgroup 13-07A   | 3                 |
| Celtuce   | 4.0               |
| Cherry subgroup 12-12A  | 0.6               |
| Corn, sweet, forage   | 9                 |
| Corn, sweet, kernel plus cob with husks removed                     | 0.4               |
| Corn, sweet, stover   | 20                |
| Cotton, gin byproducts  | 6.0               |
| Cotton, hulls   | 2.0               |
| Cotton, meal  | 1.0               |

|   |      |
|---|------|
| Cottonseed subgroup 20C   | 0.60 |
| Florence fennel   | 4.0  |
| Fruit, citrus, group 10-10  | 1.5  |
| Fruit, pome, group 11-10  | 0.20 |
| Hop, dried cones  | 20.0 |
| Kohlrabi  | 1.5  |
| Leaf petiole vegetable subgroup 22B                                 | 4.0  |
| Leafy greens subgroup 4-16A, except spinach                         | 8    |
| Peach subgroup 12-12B   | 1.5  |
| Nut, tree, group 14-12 except pistachio                             | 0.15 |
| Pepper/Eggplant, subgroup 8-10B                                     | 3.0  |
| Peppermint, tops  | 7.0  |
| Pistachio   | 0.60 |
| Plum subgroup 12-12C  | 0.6  |
| Pomegranate   | 0.5  |
| Potato, granules/flakes   | 0.40 |
| Prickly pear, fruit   | 2    |
| Prickly pear, pads  | 3    |
| Radish, tops  | 20   |
| Rapeseed subgroup 20A   | 1.5  |
| Small fruit vine climbing (except fuzzy kiwifruit), subgroup 13-07F | 3    |
| Spearmint, tops   | 7.0  |
| Spinach   | 9.0  |
| Sunflower subgroup 20B  | 0.70 |
| Tea 1   | 40   |
| Tomato, paste   | 2.0  |
| Tomato, puree   | 0.50 |
| Tomato subgroup 8-10A   | 0.4  |

|   |      |
|---|------|
| Vegetable, <i>brassica</i> , head and stem, group 5-16                        | 1.5  |
| Vegetable, cucurbit, group 9  | 1.5  |
| Vegetable, legume, bean, edible podded, subgroup 6-22A                        | 4    |
| Vegetable, legume, bean, succulent shelled, subgroup 6-22C                    | 7    |
| Vegetable, legume, pea, edible podded, subgroup 6-22B                         | 4    |
| Vegetable, legume, pea, succulent shelled, subgroup 6-22D                     | 7    |
| Vegetable, legume, pulse, bean, dried shelled, except soybean, subgroup 6-22E | 3    |
| Vegetable, legume, pulse, pea, dried shelled, subgroup 6-22F                  | 3    |
| Vegetable, root, except sugar beet, subgroup 1B                               | 0.60 |
| Vegetable, tuberous and corm, subgroup 1C                                     | 0.20 |

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)