

40 C.F.R. § 180.627

Fluopicolide; tolerances for residues.

(a) *General.* Tolerances are established for residues of the fungicide fluopicolide [2,6-dichloro-*N*-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl]benzamide], including its metabolites and degradates, in or on the commodities in the table in this paragraph. Compliance with the tolerance levels specified below is to be determined by measuring only fluopicolide [2,6-dichloro-*N*-[[3-chloro-5-(trifluoromethyl)-2-pyridinyl]methyl]benzamide] in or on the commodity.

Commodity	Parts per million
Basil, dried, leaves	200
Basil, fresh leaves	40
Bean, moth, succulent	0.90
Bean, runner, succulent	0.90
Bean, snap, succulent	0.90
Bean, wax, succulent	0.90
Bean, yardlong, succulent	0.90
Brassica, head and stem, subgroup 5A	5.0
Citrus, dried pulp	0.03
Citrus, oil	1.0
Fruit, citrus, crop group 10-10	0.01
Fruit, small, vine climbing, except fuzzy kiwifruit, subgroup 13-07F	2.0
Grape, raisin	6.0
Hop, dried cones	15
Potato, granules/flakes	0.15
Potato, processed potato waste	0.2
Vegetable, brassica (cole) leafy subgroup 5B	18

Vegetable, bulb, crop group 3–07	7.0
Vegetable, cucurbit, group 9	0.50
Vegetable, fruiting, crop group 8–10	1.6
Vegetable, leafy, except brassica, group 4	25
Vegetable, leaves of root and tuber, group 2	15.0
Vegetable root, subgroup 1A	0.15
Vegetable, tuberous and corm, subgroup 1C	0.09

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)