

40 C.F.R. § 180.680

Fluensulfone; tolerances for residues.

(a) *General.* Tolerances are established for residues of the nematicide fluensulfone, including its metabolites and degradates, in or on the commodities in the table 1 to § 180.680. Compliance with the tolerance levels specified in the following table below is to be determined by measuring only the sum of fluensulfone, 5-chloro-2-[(3,4,4-trifluoro-3-buten-1-yl)sulfonyl]thiazole and its metabolite, 3,4,4-trifluoro-but-3-ene-1-sulfonic acid, calculated as the stoichiometric equivalent of fluensulfone, in or on the commodity.

Table 1 to § 180.680

Commodity	Parts per million
Almond, hulls	5
Berry, low growing, subgroup 13-07G	0.5
Brassica, head and stem, subgroup 5A	1.5
Brassica, leafy greens, subgroup 4-16B	20
Celtuce	4
Fennel, Florence, fresh leaves and stalk	4
Fruit, citrus, group 10-10	0.3
Fruit, citrus, group 10-10, dried pulp	0.9
Fruit, citrus, group 10-10, oil	15
Fruit, pome, group 11-10	0.4
Fruit, small, vine climbing, subgroup 13-07D	0.8
Fruit, stone, group 12-12	0.15
Grape, raisin	1.5
Kohlrabi	1.5
Leaf petiole vegetable subgroup 22B	2
Leafy greens subgroup 4-16A	4

Nut, tree, group 14-12	0.02
Potato, chips	2
Potato, granules/flakes	2
Soybean, seed	0.07
Sugarcane, cane	0.06
Sugarcane, molasses	0.3
Tomato, paste	1.5
Vegetable, <i>Brassica</i> , head and stem, group 5-16	1.5
Vegetables, cucurbits, group 9	0.7
Vegetables, fruiting, group 8-10	0.7
Vegetables, leafy, except <i>Brassica</i> , group 4 ¹	4
Vegetables, leaves of root and tuber, group 2, except sugar beet	50
Vegetables, root, except sugar beet, subgroup 1B	4
Vegetables, tuberous and corm, subgroup 1C	0.8

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)