

40 C.F.R. § 63.2840

What emission requirements must I meet?

For each facility meeting the applicability criteria in § 63.2832, you must comply with either the requirements specified in paragraphs (a) through (d), or the requirements in paragraph (e) of this section. You must also comply with the requirements in paragraph (g) of this section. You must comply with the work practice standard provided in paragraph (h) of this section, if you choose to operate your source under an initial startup period subject to § 63.2850(c)(2) or (d)(2).

(a)

(1) The emission requirements limit the number of gallons of HAP lost per ton of listed oilseeds processed. For each operating month, as defined in § 63.2872, you must calculate a compliance ratio which compares your actual HAP loss to your allowable HAP loss for the previous 12 operating months as shown in Equation 1 of this section. Equation 1 of this section follows:

$$\text{Compliance Ratio} = \frac{\text{Actual Hap Loss}}{\text{Allowable Hap Loss}} \quad (\text{Eq. 1})$$

(2) Equation 1 of this section can also be expressed as a function of total solvent loss as shown in Equation 2 of this section. Equation 2 of this section follows:

$$\text{Compliance Ratio} = \frac{f * \text{Actual Solvent Loss}}{0.64 * \sum_{i=1}^n ((\text{Oilseed})_i * (\text{SLF})_i)} \quad (\text{Eq. 2})$$

Where:

f = The weighted average volume fraction of HAP in solvent received during the previous 12 operating months, as determined in § 63.2854, dimensionless. 0.64 = The average volume fraction of HAP in solvent in the baseline performance data, dimensionless. Actual Solvent Loss = Gallons of actual solvent loss during previous 12 operating months, as determined in § 63.2853. Oilseed = Tons of each oilseed type “i” processed during the previous 12 operating months, as shown in § 63.2855. SLF = The corresponding solvent loss factor (gal/ton) for oilseed “i” listed in Table 1 of this section, as follows:

Table 1 of § 63.2840—Oilseed Solvent Loss Factors for Determining Allowable HAP Loss

Type of oilseed process	A source that...	Oilseed solvent loss factor (gal/ton)

		Existing sources	New sources
(i) Corn Germ, Wet Milling	processes corn germ that has been separated from other corn components using a “wet” process of centrifuging a slurry steeped in a dilute sulfurous acid solution	0.4	0.3
(ii) Corn Germ, Dry Milling	processes corn germ that has been separated from the other corn components using a “dry” process of mechanical chafing and air sifting	0.7	0.7
(iii) Cottonseed, Large	processes 120,000 tons or more of a combination of cottonseed and other listed oilseeds during all normal operating periods in a 12 operating month period	0.5	0.4
(iv) Cottonseed, Small	processes less than 120,000 tons of a combination of cottonseed and other listed oilseeds during all normal operating periods in a 12 operating month period	0.7	0.4
(v) Flax	processes flax	0.6	0.6
(vi) Peanuts	processes peanuts	1.2	0.7
(vii) Rapeseed	processes rapeseed	0.7	0.3
(viii) Safflower	processes safflower	0.7	0.7
(ix) Soybean, Conventional	uses a conventional style desolventizer to produce crude soybean oil products and soybean animal feed products	0.2	0.2
(x) Soybean, Specialty	uses a special style desolventizer to produce soybean meal products for human and animal consumption	1.7	1.5
(xi) Soybean, Combination Plant with Low Specialty Production	processes soybeans in both specialty and conventional desolventizers and the quantity of soybeans processed in specialty desolventizers during normal operating periods is less than 3.3 percent of total soybeans processed during all normal operating periods in a 12 operating month period. The corresponding solvent loss factor is an overall value and applies to the total quantity of soybeans processed.	0.25	0.25
(xii) Sunflower	processes sunflower	0.4	0.3

This document is only available to subscribers. Please log in or purchase access.

[Purchase Login](#)